

# From the Cow's Perspective: Two Decades of Management Research at Miner Institute

**Rick Grant<sup>1</sup>**

*Member, Board of Trustees*

*William H. Miner Agricultural Research Institute*

## Summary

Two decades of management research at Miner Institute began with an understanding of behavioral time budgets - focused on eating, resting, and ruminating - and the importance of *accommodating natural cow behavior* to optimize cow comfort economics. This initial work rested squarely on the foundational work of Dr. Jack Albright at Purdue University. Twenty years ago, overcrowding was already becoming a substantial challenge for the dairy industry, and we spent considerable time assessing the behavioral, health, and production consequences of overcrowding *from the cow's perspective*. The behavioral response of cows to a given stocking density of stalls and(or) feed bunks varies by parity and lameness status, among other factors. Initial work also suggests that cows have preferences for certain stalls and(or) areas of the feed bunk and do not use pen resources uniformly. Taken together, the data strongly suggest that the stocking density experienced by an individual cow may be greater than the calculated stocking density. Stocking density should be considered a sub-clinical stressor that requires the expenditure of biological reserves, even if commonly measured performance responses remain unchanged. Additional stressors result in losses in performance and health. We have found that overcrowding of the stalls and feed bunk has a greater negative impact on rumen pH than dietary content of

physically effective fiber. The greatest duration of sub-acute rumen acidosis occurs when cows are fed a highly rumen fermentable diet, are overcrowded, and are feed restricted. Although overcrowding does not consistently reduce rumination time, it does constrain recumbent rumination, and this rumination while lying down is associated with higher rumen pH, greater intake, and percentages of milk fat and protein. Herds producing the greatest amount of de novo fatty acids and milk fat and protein manage their herds with lower feed bunk and stall stocking densities and feed rations more frequently. Ultimately, our goal has been to incorporate this behavioral management research into nutrition models, and initial, simple models have been developed and used in the field. *Time budget analysis should become an integral part of ration formulation*. Finally, at the intersection of forage quality and natural cow behavior lies the concept of precision chewing management. Optimizing management and forage-fiber characteristics so that the cow can achieve the correct balance between eating time and time for recumbent rumination appears to be central to cow health and productivity.

## Introduction

Much of this paper has been published previously in the proceedings of the Cornell Nutrition Conference (Grant, 2023). The management research at Miner Institute began

---

<sup>1</sup>Contact at: 8679 Genova Ct., Naples, FL 34114, (518) 569-0925, Email: grant@whminer.com.



with understanding behavioral time budgets and defining the concept of “cow comfort economics.” An essential insight was that we must accommodate the dairy cow’s natural behaviors – eating, resting, and ruminating – within an optimized management environment to enhance productivity, well-being, and herd profitability. Nutrition models have improved rapidly as new biology is incorporated. But the cow tells us that poor management constrains her ability to respond to the diet. Work by Bach et al. (2008) quantified the overriding impact of non-dietary factors, such as feed availability and stall stocking density, on farm-to-farm variability in milk production relative to nutritional factors. As a result of this insight, a long-term goal of Miner research has been to incorporate cow management into nutrition models for ration formulation.

I’ve attempted to summarize what we have learned over two decades about optimal management from the cow’s perspective: cows that aren’t rushed while eating, have the freedom to lie down and ruminate, and can strike the correct balance between eating and recumbent rumination, will have optimal rumen conditions for fiber digestion, microbial growth, and healthy production of more milk components. Explaining how I arrived at this statement is the objective of this paper.

### **Time Budgets and Cow Comfort Economics**

We routinely monitor time budget behaviors to assess the quality of herd management. But what is commonplace today was not in the 1970s and earlier, and we can thank Dr. Jack Albright of Purdue University for his pioneering insight into the value of studying cow behavior to improve performance and well-being. In the June 2023 issue of the Miner Institute Farm Report (Grant, 2023), I explained how Albright and his team in 1975

recorded the daily behavior of the first cow to break 22,676 kg of milk/yr (i.e., 50,000 lb/yr), Beecher Arlinda Ellen. This record-setting cow spent approximately 6 hr/day eating and nearly 14 hr/day lying down. Her daily dry matter intake (**DMI**) approached an enormous 7% of body weight. Ellen spent 7.5 hr/day ruminating with about 93% of that rumination occurring while lying down (i.e., while sternally recumbent). Albright suggested that sternal recumbency, with a slight left-side laterality, allowed the cow to remain comfortable and for the rumen to function optimally. Given Ellen’s prodigious productivity, it is unsurprising that recently published research has reported a positive relationship between recumbent rumination and feed intake, and milk fat and protein percentages (McWilliams et al., 2022). More on this relationship later.

Conversations with Albright in the mid- to late 1990s led to an active interest in assessing cow behavior on commercial dairy farms. Baseline behavioral data collected by Bill Matzke (2003) as part of his MS research at the University of Nebraska on commercial dairy farms helped develop realistic and useful time budgets for lactating cows. Insights from this early work on time budgeting were: 1) excessive time outside the pen has a profound negative impact on resting behavior; 2) cows locked in headlocks beyond 1 hr stop eating, and although 30 to 40% begin to ruminate while standing, they are essentially wasting their time; 3) the highest milk-producing cows in a pen spend up to 14 hr/day resting (as did Beecher Arlinda Ellen); and 4) cows forced by poor management to spend more time on non-eating activities borrow that time from another one – most often resting. This early work highlighted the natural relationship between eating and resting, and how this interplay profoundly affects production and health.

Research at Miner Institute in the early 2000s cemented the idea that eating, resting, and ruminating are linked biologically – forming the foundation of cow comfort and what we called cow comfort economics (Grant, 2003; 2004; 2015). In many ways, we rediscovered previously reported information on the connection between eating and resting, and the priority that cows place on resting versus eating behavior. Time outside the pen was crucial, given its impact on time available for eating, resting and other behaviors, and was viewed as the most vital component of dairy cow well-being in a competitive free-stall environment. Time outside the pen literally sets the limit on what is possible behaviorally within the pen. An Excel spreadsheet (Time Budget Evaluator, version 3. [www.whminer.org](http://www.whminer.org); described by Grant, 2004) was developed to assess the time budget behaviors of cows and the potential for lost production in competitive environments.

### **Stocking Density and Overcrowding from the Cow's Perspective**

Twenty years ago, it was already clear that overcrowding was becoming a significant management challenge in the dairy industry. When a pen of cows is overstocked, the availability of resting and feeding space, area per cow, and access to water are all restricted. Studies conducted in the past 20 years have amply demonstrated that overcrowding, especially beyond 120% of stalls, hampers lying time, boosts lameness, reduces milk yield, encourages undesirable feeding behaviors (such as slug feeding and sorting), and elevates somatic cells (reviewed by Grant, 2007; 2015). Early on, we compared alternative models for studying stocking density and concluded that simply denying resting and feeding space to simulate overcrowded 4-row housing was bioequivalent to more complicated research models (Krawczel et al., 2012b).

Defining the differential effect of overcrowding in a 4- or 6-row barn has not been rigorously examined with controlled research, yet we can safely infer that any specific stall stocking density will have potentially greater negative effects in a 6- versus a 4-row barn. Importantly, we have proposed that *time outside the pen interacts with stocking density*. Even at 100% stall stocking in a 4-row barn, time outside the pen of 6 versus 3 hr/day reduced resting time by 2.6 hr/day for multiparous cows and 4.2 hr/day for primiparous cows in commingled pens (Matzke, 2003). It is difficult to name many factors that would affect resting so substantially.

### *Parity and lameness affect response to overcrowding*

Soon after my arrival at Miner Institute, Chris Hill began a series of studies aimed at defining the short-term effects of overcrowding on behavior, performance, and how the response differed by parity and lameness. Hill et al. (2006; 2009) homed in on how stocking density affected primi- versus multiparous cows. As stall and headlock stocking densities in 2-row pens increased from 100 to 142%, the difference in milk yield between multi- and primiparous cows increased from 2.8 kg/day at 100% up to 9.6 kg/day for 130% stocking density. At 142% stocking density, the difference dropped to 6.8 kg/day, reflecting a negative effect on the dominant, multiparous cows as well as the subordinate, primiparous cows at this elevated level. Similarly, the difference in milk yield between sound and lame cows increased by 11.9 kg/day as stocking density of stalls and headlocks increased. Reinforcing the importance of resting/rumination, these reductions in milk yield tracked with losses in both lying and ruminating, particularly for the lame cows.

### *What stocking density does the cow actually experience?*

In this early work, we observed that sometimes daily rumination time was affected by stocking density, but not always (Hill, 2006; Krawczel et al., 2012a). But in all our studies, rumination while lying down decreased at higher stocking densities. Furthermore, one study noted that subordinate cows lying in stalls preferred by dominant cows spent 40% less time ruminating while lying in the stall (Krawczel, 2007, unpublished). Given the recently recognized importance of recumbent rumination on DMI and milk components, this reduction in resting rumination with higher stocking density looms large as an economically important management challenge. An overarching management question must be: *what stocking density does an individual cow experience relative to the measured value (e.g., cows/stall)?* Beyond the number of cows per stall, it seems likely that subordinate cows faced with using stalls preferred by dominant cows must experience a functionally higher stocking density than the calculated one.

Currently, we are evaluating the importance of location preference within a pen, whether free stalls, alley, or feed bunk. If individual cows, or types of cows, prefer to eat or rest only in certain areas of a pen, then that would also affect their perceived level of competition for the resource and the optimal management practice to ensure adequate access for all cows in the pen. For example, Hefter et al. (2023) reported that lame cows preferred to use the free stalls nearest the pen exit. Cows, regardless of lameness status, also exhibited a strong preference for eating at feed bunk sections closest to the exit gate between approximately 6:00 am and 9:00 pm, but not at night, presumably in relation to milking and feed availability. Most recently, Pupo et al. (2024) found that approximately 25% of cows

in a multi-study dataset had predictable feed-bunk location preference and about 1 kg/day lower DMI. Overall, location preference within a pen deserves more research given its evident fundamental importance to precise, individual cow management.

De Vries et al. (2016) published a model that predicts changes in performance and profit per stall with variable stall stocking densities. A central relationship in this model of a loss of 0.50 kg/day of milk per 10% greater stall stocking density is based on Grant (2011), Fregonesi et al. (2007), and Bach et al. (2008). This model is useful to project how stocking density affects specific pens and herds of cows and is available on the University of Florida web site: <https://edis.ifas.ufl.edu/publication/AN346>.

### *Assessing cow comfort in overcrowded pens*

Increasing stocking density beyond 100 to 113% with our research model clearly increases the proportion of cows standing in alleys and compromises their ability to access free stalls when motivation to lie down is greatest. Pete Krawczel led research demonstrating that assessing cow comfort on a pen level is best done using the stall use index which is calculated as the number of cows in a pen lying down divided by the total number of cows minus those actively eating (Krawczel et al., 2008). In other words, the denominator includes cows that are essentially wasting their time, idling and mainly waiting to use a stall. The commonly used cow comfort index and stall standing index, calculated using only the cows lying or standing in a stall, remain relatively unchanged at higher stocking densities. For indices routinely measured and monitored on-farm, the stall use index and the rumination index (% of cows ruminating that are lying in stalls) top our list of priority indices. Campbell (2017) found that the commonly used rumination index is related to 24-hr rumination

time. Importantly, the rumination index allows us to monitor recumbent rumination specifically.

### **Feed and Feeding Environment: Focus on Rumen pH**

#### *Overcrowding as a subclinical stressor*

Given the wide array of negative behavioral, health, and performance consequences of overcrowding measured in research studies, the diversity of herd responses to overcrowding seems puzzling. Some herds seem to be immune to the negative consequences of overcrowding, whereas others experience severe effects at low stall stocking density, and everything in between. One possible explanation would be to consider overcrowding a subclinical stressor. Moberg (2000) defined a subclinical stressor as one that depletes biological resources of an animal without creating detectable change in function, such as milk yield, reproduction, or health. However, it leaves the animal unable to successfully respond to additional stressors. Using this model, one can imagine an overcrowded pen of dairy cattle that appears to be meeting most or even all the commonly used industry performance benchmarks, but at some point, secondary stressor(s) will cause a measurable change in function. We propose that the extent to which the biological reserves are expended by the subclinical stress of overcrowding in any specific herd is a function of the quality of the housing and management routines.

Mac Campbell embraced this model and asked the question of what occurs with chronically overcrowded dairy cows when the secondary stressors were inadequate ration fiber and restricted feed availability (Campbell and Grant, 2016). As a ruminant, little is as important to dairy cattle well-being and productivity as a healthy rumen environment, particularly pH.

When comparing stall and headlock stocking density of 100 versus 142% with physically effective neutral detergent fiber (**peNDF**) of 19 or 21% (8.5 and 9.7% undegraded NDF at 240 hr of in vitro fermentation; uNDF<sub>240</sub>), Campbell and Grant (2016) reported that the extent of subacute rumen acidosis (time that pH < 5.8) was up to 2 hr/day greater for the main effect of overstocking versus fiber. Furthermore, when cows were fed the same total mixed ration (**TMR**), cows overstocked at 142% and also experiencing a 5-hr feed restriction prior to delivery of the morning ration had up to 9 hr/day greater subacute rumen acidosis than those at 100% stocking density with unrestricted feed access. The bottom line was that stocking density and feed restriction had a greater negative impact on rumen pH than dietary fiber characteristics. In fact, this work suggests that the ideal approach for lowering rumen pH would be to: 1) feed a highly fermentable diet, 2) overcrowd the feed bunk and stalls, and 3) feed to an empty bunk.

Although this work does not give the nutritionist and formulated ration a pass when it comes to rumen pH, it clearly recasts the relative importance of ration formulation and feed-bunk management. *Ration formulation, especially carbohydrates, is critical to maintaining healthy rumen pH conditions, but we now know that how the ration is fed is paramount.*

Overcrowding resulted in lower rumen pH in Campbell's work, but it did not have consistent effects on feeding or ruminating time. Like previous research, lying time was significantly reduced, and notably, recumbent rumination was depressed at higher stocking density. With overcrowded environments, Campbell (2017) observed a negative relationship between the fraction of rumination that occurs in the free stall (x) and the hours that rumen pH is below 5.8 ( $y = -20.7x + 21.1$ ;  $r = 0.66$ ). This result had not been expected but raised important

questions regarding the potential importance of rumination in stalls to maintaining desirable rumen pH conditions. For the first time, *there was evidence that posture during rumination (i.e., standing or lying) affected rumen pH*. This makes sense given what we know about the relative rates of saliva secretion when the cow is eating, ruminating, or resting, and how sternal recumbency aids the rumination process (reviewed by Grant and Cotanch, 2023).

## Management and Milk Components

Based on the positive relationship between rumen pH and greater milk de novo fatty acids and total fat production (Fukumori et al., 2021), de novo fatty acid content of milk fat should serve as a useful barometer of rumen pH conditions conducive to fiber digestion and microbial growth. Barbano has published extensively on the positive relationship between milk de novo fatty acids and output of milk fat and true protein (Barbano et al., 2018). Published literature demonstrates the positive relationship between higher rumen pH and greater de novo fatty acids, and milk fat, and true protein percentages (Allen, 1997; Fukumori et al., 2021; Stone, 2004).

At roughly the same time that Campbell was conducting his work, Melissa Woolpert reported on factors affecting milk de novo fatty acid production for 79 commercial herds in VT and NY (Woolpert et al., 2016; 2017). Higher de novo fatty acid herds produced 17% more milk fat and 14% more true protein than lower de novo herds. Across these herds, the top-five factors that characterized high de novo fatty acid, high milk component herds were: 1) dietary fat  $\leq 3.5\%$  of ration DM; 2) dietary peNDF  $\geq 21\%$  of ration DM; 3 and 4) lower feed bunk and stall stocking densities; and 5) greater feeding frequency of TMR. High de novo herds were 10 times more likely than low de novo herds to

have feed bunk space  $\geq 45$  cm/cow (18 in/cow) and stall stocking density  $\leq 110\%$ . High de novo herds were 5 times more likely to feed TMR twice daily rather than once. Importantly, 65% of the variation in milk de novo fatty acid content among herds was explained by feed bunk space alone. This relates well to earlier work that found greater feed bunk space to be correlated with increased milk yield and milk fat percentage (Sova et al., 2013). We cannot overstate the negative effect that overcrowding may have on milk component production. *Overcrowded cows cannot respond maximally in milk components to the formulated ration.*

Considering Campbell's demonstration that greater rumination in the stalls is associated with higher rumen pH, then cows in competitive feeding environments who can achieve more recumbent rumination should have better rumen conditions for fiber digestion and potential for greater de novo fatty acids, and milk fat and protein outputs. In fact, McWilliams et al. (2022) observed that cows with greater ruminating time while lying down had greater DMI and produced milk with more fat and true protein contents. To follow up on this work, we conducted a student research project in spring 2023 (unpublished) using Holstein cows that ranged from 3.2 to 6.4% milk fat. Like McWilliams et al. (2022), we found a significant positive correlation ( $R = 0.34$ ;  $P = 0.03$ ) between minutes of rumination while lying down and milk fat percentage. This relationship was significantly stronger than for ruminating, resting, or eating time. Recumbent rumination, and not simply total *rumination time, is critical to a healthy, productive, and profitable herd.*

## Precision Chewing Management: Forage Quality and the Balance Between Eating and Recumbent Rumination

Eating time between 3 and 5 hr/day is typically associated with desirable feeding

behavior as reviewed by Grant and Albright (2001). In a cow's ideal environment, she will achieve over 80% of daily rumination while lying down. Hence, the cost of excessive time at the bunk, beyond 5 hr/day, is considerable since it directly reduces time available for resting and ruminating. In fact, Jiang et al. (2017) observed an exact daily balance between total chewing time, driven by eating, and resting time. As dietary forage content increased beyond 50% of ration DM, eating time increased markedly, with minor impact on rumination. The extra time needed for eating was carved minute-for-minute from resting time, and inescapably from resting rumination. *We propose that maintaining a proper balance between eating time (3 to 5 hr/day) and optimal resting (11 to 14 hr/day) and ruminating time (8 to 9 hr/day, >80% while lying) is central to cow productivity and well-being.*

Understanding the fundamental importance of maintaining the balance between eating and recumbent rumination was paradigm shifting. And, it begged the question of how forage quality influences the cow's ability to keep eating and resting/ruminating in balance. Fiber plays a key role in stimulating chewing, both eating and ruminating. Dietary fiber content, source of fiber, particle size, digestibility, and fragility all contribute to the effect of fiber on chewing. Grant and Cotanch (2023) reviewed recent research showing that the primary effect of forage particle size is on eating rather than ruminating in many feeding scenarios. Cows chew to a relatively uniform particle size endpoint prior to swallowing while eating. Consequently, the rumen is populated with a forage particle size distribution that is smaller and more uniform than the TMR. In fact, for many of our commonly fed corn silage-based diets, the particles retained on the 8-mm sieve of the Penn State Particle Separator are essentially the same size as the swallowed bolus while eating.

Particle size reduction that occurs during eating requires more chews per gram of feed DM and takes longer for diets that are coarser, higher in forage NDF, and less digestible. We have measured up to a 6-fold reduction in the longest TMR particles prior to ingestive swallowing when we fed cows coarse diets with high uNDF240 content (Grant et al., 2018; Smith, 2019). The published literature shows that eating time can be increased by up to an hour when forage NDF is greater, less digestible, or chopped coarser (Grant and Ferraretto, 2018; Grant and Cotanch, 2023). Details are provided in the review by Grant and Cotanch (2023), but the main point is that a practical, on-farm system has been proposed that allows theoretical length of cut to be adjusted based on forage maturity, moisture content, digestibility, and(or) fragility to optimize the balance between eating and recumbent rumination. Additionally, an optimal TMR particle size distribution has been proposed to balance eating, ruminating, and resting.

In our approach to particle size, *eating time at the feed bunk is a crucial and overlooked component of forage quality*. Lower quality forage is less digestible and takes longer to eat unless particle size is reduced. Likewise, higher quality forage benefits from longer chop length in most cases. Understanding the fundamental importance of this relationship will be most critical in competitive feeding environments – which characterize many of our commercial herds. We view this as a holistic approach to optimizing the two fundamental pillars of a profitable dairy farm: forage quality and cow comfort.

### **Modeling Management: The Holy Grail?**

Publications by Grant and Tylutki (2010; 2011) for the first time proposed that time budget analysis should become a routine

and important part of DMI prediction and ration formulation. The feeding environment is comprised of both physical and social components that modulate feeding behavior and feed intake in dairy cattle. Currently, key components of the physical environment, such as temperature, humidity, wind speed, and so forth, are inputted into the model during ration formulation. But, effectively incorporating social and management factors, such as time budgets, stocking density, and feed availability, has proven challenging. Nonetheless, nutrition models need to incorporate management inputs. Feeding behavior and intake are dramatically affected by time available to eat, forage NDF characteristics, notably particle size, and stocking density to name a few.

To date, the most useful effort has been the management model created as part of Michael Miller's Ph.D. dissertation (Miller et al. 2020; detailed equations provided by Miller, 2020ab). The model remains a work-in-progress, but field experience over the past few years suggests that it has usefulness. It has been implemented in AMTS version 4.18 as a recipe tool called the "management model."

The model was designed to allow input of commonly measured farm variables, such as stocking density and milking time, to assess the effect of management decisions on DMI, milk production, and behavior. The model is divided into five sections: 1) behavioral time budget adapted from the original model by Grant and Tylutki (2011); 2) stocking density calculation; 3) eating time prediction; 4) DMI prediction; and 5) physically effective uNDF240 (**peuNDF240**) adjustment to DMI. The time budget analysis provides time available for eating plus resting, and eating time is predicted from NDF, peNDF, body weight, milk yield, and feeding frequency. Once time available for rest has been calculated, it is adjusted based on stocking density of stalls

or manger, depending on whether the barn uses headlocks or post-and-rail. Then, we use a relationship between resting time and milk yield, based on a review of published data to predict fat-corrected milk (**FCM**) production. The FCM value, together with body weight and week of lactation, is used to predict DMI using the NRC (2001) equation. Although this approach backs into a DMI prediction, it seems to be the best approach for now since no reported relationship between stocking density and short-term DMI has been reported.

Three main limitations remain: 1) rigorous model validation; 2) adjustments for parity effects; and 3) incorporation of feed availability. Although we know that parity and feed availability are important, a lack of data hampers further model development. The model is most sensitive to milking time and stocking density which makes sense given how important they are on farm. As part of model development, Miller (2020) reviewed the published literature and updated equations relating stocking density with lying time and lying time with milk yield. Although not a main part of the model, an intake adjustment based on the relationship between  $peuNDF_{240}$  and DMI was also incorporated to take advantage of the database we have built comparing  $uNDF_{240}$  and  $peuNDF_{240}$  and their relationship with DMI and energy-corrected milk (Grant et al., 2018; Miller et al., 2020; Farricker et al., 2022). It is conceivable that this component in the future could also incorporate the interaction between rumen fermentable starch and  $peuNDF_{240}$  (Smith et al., 2020).

### Looking to the Future!

The amount of cattle management research conducted over the past 20 years is mind boggling. When reviewing the many management factors affecting cow performance and health, few are surprising. *What is surprising*

is the sheer magnitude of the cow responses to an improvement in comfort. Furthermore, it is amazing how a few factors rise repeatedly to the top as essential for a low-stress management environment: 1) time available for eating, resting, and ruminating; 2) managing stocking density and overcrowding; 3) ensuring feed availability 24/7; and 4) resting area comfort (e.g., deep bedding). Regardless of the management system – free-stall and parlor, automated milking system, tie stall, pasture, or whatever evolves in the future – we need to understand how to accommodate natural cow behavior. Any sustainable dairy system will consider environmental, welfare, and societal concerns. And from a profitability perspective, there is little doubt that cow comfort economics will remain compelling.

Research needs to continue that builds on our new understanding of forage quality and cow well-being. How forage characteristics, housing, and management environment affect each cow's ability to balance eating with recumbent rumination needs to be a focus going forward. We have termed this concept *Precision Chewing Management*. As advances in nutrition models occur, we need to implement a functional model incorporating physical and social components of the cow's environment. Time budgeting should become a routine part of ration formulation.

As behavior research moves toward answering much-needed questions that cross-cut society, such as cow-calf separation, we cannot forsake research aimed at enhancing the productivity and well-being of cows in commercial management systems. So, we return to the original assertion of this paper that I believe captures the essential importance of our work at Miner Institute: “cows that aren't rushed while eating, have the freedom to lie down and ruminate, and can strike the correct balance between eating and recumbent

rumination, will have optimal rumen conditions for fiber digestion, microbial growth, and healthy production of more milk components.”

Our management research at Miner Institute reaches back to Jack Albright's pioneering work in cattle behavior; his recognition that the need to accommodate natural cow behavior would become an essential component of successful herd management. After briefly reviewing 2 decades of management research at Miner and looking to the future, I conclude that applied dairy management research must continue with the goal of “unleashing every cow's inner Ellen!”

## References

- Allen, M.S. 1997. Relationship between fermentation acid production in the rumen and the requirement for physically effective fiber. *J. Dairy Sci.* 80:1447–1462.
- Bach, A., N. Valls, A. Sloans, and T. Torrent. 2008. Associations between non-dietary factors and dairy herd performance. *J. Dairy Sci.* 91:3259-3267.
- Barbano, D.M., H. Dann, C. Melilli, and R. Grant. 2018. Milk analysis for dairy herd management: Today and in the future. Pages 103-115 in *Proc. Cornell Nutr. Conf. for Feed Manufac.* East Syracuse, NY. Cornell University, Ithaca, NY.
- Campbell, M.A. 2017. Interaction of stocking density and the feeding environment in lactating Holstein dairy cows. Ph.D. Dissertation. University of Vermont, Burlington.
- Campbell, M.A., and R.J. Grant. 2016. Interactions between the feed and feeding environment. Pages 49-58 in *Proc. Cornell Nutr. Conf. for Feed Manufac.* East Syracuse, NY. Cornell University, Ithaca, NY.

- De Vries, A., H. Dechassa, and H. Hogeveen. 2016. Economic evaluation of stall stocking density of lactating dairy cows. *J. Dairy Sci.* 99:3848-3857.
- Farricker, M., J. Darrah, A. Pape, M. Miller, and R. Grant. 2022. Ration content of undegradable and physically effective neutral detergent fiber and its relationship with dry matter intake and energy-corrected milk yield of Holstein cows. *J. Dairy Sci.* 105 (Suppl. 1):246 (Abstr.).
- Fregonesi, J.A., C.B. Tucker, and D.M. Weary. 2007. Overstocking reduces lying time in dairy cows. *J. Dairy Sci.* 90:3349-3354.
- Fukumori, R., W. Shi, S. Oikawa, and M. Oba. 2021. Evaluation of relationship between ruminal pH and the proportion of de novo fatty acids in milk. *JDS Comm.* 2:123-126.
- Grant, R. J. 2003. Taking advantage of dairy cow behavior: Cost of ignoring time budgets. Pages 67-76 in *Proc. Cornell Nutr. Conf. for Feed Manufac.* East Syracuse, NY. Cornell University, Ithaca, NY.
- Grant, R.J. 2004. Incorporating dairy cow behavior into management tools. Pages 65-76 in *Proc. Cornell Nutr. Conf. for Feed Manufac.* East Syracuse, NY. Cornell University, Ithaca, NY.
- Grant, R.J. 2007. Effect of stocking density on cow comfort, health, and productivity. Pages 105-111 in *Proc. Cornell Nutr. Conf. for Feed Manufac.* East Syracuse, NY. Cornell University, Ithaca, NY.
- Grant, R. 2011. Taking advantage of natural behavior improves dairy cow performance. eXtension. <http://www.extension.org/pages/11129/taking-advantage-of-natural-behavior-improves-dairy-cow-performance>.
- Grant, R.J. 2015. Economic benefits of improved cow comfort. Publ. by Novus Int., Inc. St. Charles, MO Access on line: [https://www.dairychallenge.org/pdfs/2015\\_National\\_resources/Novus\\_Economic\\_Benefits\\_of\\_Improved\\_Cow\\_Comfort\\_April\\_2015.pdf](https://www.dairychallenge.org/pdfs/2015_National_resources/Novus_Economic_Benefits_of_Improved_Cow_Comfort_April_2015.pdf)
- Grant, R.J. 2023. Diary of a world record holder. The William H. Miner Agric. Res. Inst. Farm Report. June 2023. Page 1.
- Grant, R.J. 2023. What we've learned from cows: A tale of two decades of management research at Miner. *Proc. Cornell Nutr. Conf. Feed Manufac.* East Syracuse, NY. Cornell University, Ithaca, NY. <https://hdl.handle.net/1813/36419>
- Grant, R.J., and J.L. Albright. 2001. Effect of animal grouping on feeding behavior and intake of dairy cattle. *J. Dairy Sci.* 84(E. Suppl.):E156-E163.
- Grant, R.J., and K.W. Cotanch. 2023. Chewing behavior of dairy cows: Practical perspectives on forage fiber and the management environment. *Appl. Anim. Sci.* 39:146-155.
- Grant, R.J., and L.F. Ferraretto. 2018. Silage review: Silage feeding management: Silage characteristics and dairy cow feeding behavior. *J. Dairy Sci.* 101:4111-4121.
- Grant, R.J., W.A. Smith, M.D. Miller, K. Ishida, and A. Obata. 2018. Relationships between undigested and physically effective fiber in lactating dairy cows. *Proc. Cornell Nutr. Conf. for Feed Manufac.* East Syracuse, NY. Cornell University, Ithaca, NY.
- Grant, R.J., and T. Tylutki. 2011. Influence of social environment on feed intake of dairy cattle. Pages 86-100 in *Proc. Cornell Nutr. Conf. for Feed Manufac.* East Syracuse, NY. Cornell University, Ithaca, NY.

- Grant, R.J., T. Tylutki, and P. Krawczel. 2010. Modeling the influence of management on dry matter intake in dairy cattle. *J. Dairy Sci.* 93 (E-Suppl 1):468. (Abstr.).
- Hefter, K.R., J.G. Owen, A.E. Pape, S.Y. Morrison, and R.J. Grant. 2023. Effect of lameness on location preference within a pen for lactating dairy cows. *J. Dairy Sci.* 106(Suppl. 1):307 (Abstr.).
- Hill, C.T. 2006. The effects of stocking rate, parity, and lameness on the short-term behavior of dairy cattle. M.S. Thesis. University of Vermont, Burlington.
- Hill, C.T., P.D. Krawczel, H.M. Dann, C.S. Ballard, R.C. Hovey, W.A. Falls, and R.J. Grant. 2009. Effect of stocking density on the short-term behavioural responses of dairy cows. *App. Anim. Behav. Sci.* 117:144-149.
- Jiang, F.G., X.Y. Lin, Z.G. Yan, Z.Y. Hu, G.M. Liu, Y.D. Sun, X.W. Liu, and Z.H. Wang. 2017. Effect of dietary roughage level on chewing activity, ruminal pH, and saliva secretion in lactating Holstein cows. *J. Dairy Sci.* 100:2660-2671.
- Krawczel, P.D., C.T. Hill, H.M. Dann, and R.J. Grant. 2008. Short communication: Effect of stocking density on indices of cow comfort. *J. Dairy Sci.* 91:1903-1907.
- Krawczel, P.D., L.B. Klaiber, R.E. Butzler, L.M. Klaiber, H.M. Dann, C.S. Mooney, and R.J. Grant. 2012a. Short-term increases in stocking density affect the lying and social behavior, but not the productivity, of lactating Holstein dairy cows. *J. Dairy Sci.* 95:4298-4308.
- Krawczel, P.D., C.S. Mooney, H.M. Dann, M.P. Carter, R.E. Butzler, C.S. Ballard, and R.J. Grant. 2012b. Effect of alternative models for increasing stocking density on the short-term behavior and hygiene of Holstein dairy cows. *J. Dairy Sci.* 95:2467-75.
- Matzke, W.C. 2003. Behavior of large groups of lactating dairy cattle housed in a free stall barn. M.S. Thesis. University of Nebraska, Lincoln.
- McWilliams, C.J., A.J. Schwanke, and T.J. DeVries. 2022. Is greater milk production associated with cows who have a greater probability of ruminating while lying down? *JDS Comm.* 3:66-71.
- Miller, M.D. 2020. Effects of forage quality and management decisions on feed intake of lactating dairy cows. Ph.D. dissertation. University of Vermont, Burlington.
- Miller, M.D., W.A. Smith, and R.J. Grant. 2020a. Relationship of undigested and physically effective neutral detergent fiber with dry matter intake and energy-corrected milk yield of Holstein cows. *J. Dairy Sci.* 103(Suppl. 1):279(Abstr.).
- Miller, M.D., T.P. Tylutki, and R.J. Grant. 2020b. Modeling the effect of social environment on dry matter intake: Time budget behaviors and stocking density. Proc. 2020 Virtual Cornell Nutrition Conference. Cornell Univ., Ithaca, NY. <https://ecommons.cornell.edu/handle/1813/72890>
- Moberg, G.P. 2000. Biological response to stress: Implications for animal welfare. Pages 1-21 in *The Biology of Animal Stress: Basic Principle and Implications for Animal Welfare*. CABI Publishing. London, UK.

National Research Council (NRC). 2001. Nutrient requirements of dairy cattle. 7th rev. ed. Nat. Acad. Press. Washington, DC.

Pupo, M.R., J.R. R. Dorea, and L.F. Ferraretto. 2024. Determining feedbunk preference and its association with performance of dairy cows. *J. Dairy Sci.* 107 (Suppl. 1): 389 (Abstr.).

Smith, W.A. 2019. Relationships between undigested and physically effective fiber in lactating dairy cow diets. M.S. thesis. Univ. of Vermont, Burlington.

Smith, K.M., R.J. Grant, and A. Obata. 2020. Relationships between starch and physically effective and undegraded fiber in lactating dairy cows. Presented at 2020 Virtual Cornell Nutrition Conference. Cornell Univ. Ithaca, NY. <https://ecommons.cornell.edu/handle/1813/72890>

Sova, A.D., S.J. LeBlanc, B.W. McBride, and T.J. DeVries. 2013. Associations between herd-level feeding management practices, feed sorting, and milk production in freestall dairy farms. *J. Dairy Sci.* 96:1-12.

Stone, W.C. 2004. Nutritional approaches to minimize subacute ruminal acidosis and laminitis in dairy cattle. *J. Dairy Sci.* 87:(E. Suppl.): E13–E26.

Woolpert, M.E., H.M. Dann, K.W. Cotanch, C. Melilli, L.E. Chase, R.J. Grant, and D.M. Barbano. 2016. Management, nutrition, and lactation performance are related to bulk tank milk de novo fatty acid concentration on Northeastern US dairy farms. *J. Dairy Sci.* 99:8486-8497.

Woolpert, M.E., H.M. Dann, K.W. Cotanch, C. Melilli, L.E. Chase, R.J. Grant, and D.M. Barbano. 2017. Management practices, physically effective fiber, and ether extract are related to bulk tank milk de novo fatty acid concentration on Holstein dairy farms. *J. Dairy Sci.* 100:5097–5106.